

NEXN Rabbit pAb

货号: **AYP19457**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	72kDa/73kDa/79kDa/80kDa
实测分子量	81kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SGC-7901,Mouse testis,Rat brain,Rat spleen,Rat testis
细胞定位	Cell junction,Cytoplasm,Z line,adherens junction,cytoskeleton,myofibril,sarcomere
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 516-675 of human NEXN (NP_653174.3).
序列	RFEQMAKAREEEEEQRRIIEEQKLLRMQFEQREIDAALQKKREEEEEEGSIMNGSTAEDEEQTRSGAPWFKKPLKNTSVVD SEPVRFTVKVTGEPKPEITWWFEGEILQDGEDYQYIERGETYCLYLPETFPEDGGEYMCKAVNNKGSAASTCILTIESKN

靶点信息

研究背景	This gene encodes a filamentous actin-binding protein that may function in cell adhesion and migration. Mutations in this gene have been associated with dilated cardiomyopathy, also known as CMD1CC. Alternately spliced transcript variants have been described.
基因ID	91624

基因名	NEXN
Swiss	Q0ZGT2
别名	NEXN;CMH20;NELIN;nexilin

产品验证



Western blot analysis of NEXN expressed in Mouse testis,Rat brain,SGC-7901 using NEXN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn