

— ABLYBIO, Help Your Research



# SCD5 Rabbit pAb

货号: **AYP19458**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	28kDa/37kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG, 293T
细胞定位	Endoplasmic reticulum membrane, Multi-pass membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-75 of human SCD5 (N P_001032671.2).
------	---

## 靶点信息

研究背景	Stearoyl-CoA desaturase (SCD; EC 1.14.99.5) is an integral membrane protein of the endoplasmic reticulum that catalyzes the formation of monounsaturated fatty acids from saturated fatty acids. SCD may be a key regulator of energy metabolism with a role in obesity and dislipidemia. Four SCD isoforms, Scd1 through Scd4, have been identified in mouse. In contrast, only 2 SCD isoforms, SCD1 (MIM 604031) and SCD5, have been identified in human. SCD1 shares about 85% amino acid identity with all 4 mouse SCD isoforms, as well as with rat Scd1 and Scd2. In contrast, SCD5 shares limited homology with the rodent SCDs and appears to be unique to primates (Wang et al., 2005 [PubMed 15907797]).[supplied by OMIM, Mar 2008]
基因ID	79966
基因名	SCD5
Swiss	Q86SK9 ( <a href="https://www.uniprot.org/uniprotkb/Q86SK9/entry">https://www.uniprot.org/uniprotkb/Q86SK9/entry</a> )
别名	SCD5,ACOD4,FADS4,HSCD5,SCD2,SCD4,SCD5 Rabbit pAb,Acyl-CoA-desaturase 4,Stearoyl-CoA 9-desaturase,Stearoyl-CoA desaturase 2

## 产品验证



Western blot analysis of SCD5 expressed in U-87MG,293T using SCD5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)