— ABLYBIO, Help Your Research

CDK5RAP3 Rabbit pAb

货号: AYP19459

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	**WB:** 1:500 - 1:2000
理论分子量	31kDa/46kDa/56kDa/59kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T,LO2
细胞定位	Cytoplasm,Nucleus,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human CDK5R AP3 (NP_788276.1).

靶点信息

研究背景	This gene encodes a protein that has been reported to function in signaling pathways governing transcriptional regulation and cell cycle progression. It may play a role in tumorigenesis and metastasis. A pseudogene of this gene is located on the long arm of chromosome 20. Alternative splicing results in multiple transcript variants that encode different isoforms.
基因ID	80279
基因名	CDK5RAP3
Swiss	Q96JB5 (https://www.uniprot.org/uniprotkb/Q96JB5/entry)
别名	CDK5RAP3,C53,HSF-27,IC53,LZAP,MST016,OK/SW-cl.114,PP1553,CDK5RAP3 Rabbit pAb,CDK5 activator-binding protein C53,LXXLL/leucine-zipper-containing ARF-binding protein,Protein HSF-27

产品验证



Western blot analysis of CDK5RAP3 expressed in U-87MG,293T,LO2 using CDK5RAP3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)