

CWC22 Rabbit pAb

货号: **AYP19474**

产品信息

| | |
|-------|--|
| 反应 | Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:1000 - 1:2000 |
| 理论分子量 | 105kDa |
| 实测分子量 | 110kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse liver,Mouse testis |
| 细胞定位 | Nucleus,Nucleus speckle |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 639-908 of human CWC 22 (NP_065994.1). |
| 序列 | LTDELREHLKNTPKVIVAQKPDVEQNKSSPSSSSSASSSESDDSSDSDSDSSDSSSESSEESDSSSISSHSSASANDVRK KGHGKTRSKEVDKLRNQQTNDRKQKERRQEHGHQETRTERERRERSEKHRDQNSSGSNWRDPITKYTSDKDVPSENNY SRVANDRDQEMHIDLENKHGDPKPKKRGERRNSFSENEKHTRIKDSENFRRKDRSKSKEMNRKHSGSRSDDEDRYQNG AERRWEKSSRYSEQSRESKKNQDRRREKSPAKQK |

靶点信息

| | |
|------|--|
| 研究背景 | Required for pre-mRNA splicing as component of the spliceosome. Promotes exon-junction complex (EJC assembly). Hinders EIF4A3 from non-specifically binding RNA and escorts it to the splicing machinery to promote EJC assembly on mature mRNAs. Through its role in EJC assembly, required for nonsense-mediated mRNA decay. |
|------|--|

| | |
|-------|------------------------|
| 基因ID | 57703 |
| 基因名 | CWC22 |
| Swiss | Q9HCG8 |
| 别名 | CWC22;EIF4GL;NCM;fSAPb |

产品验证



Western blot analysis of CWC22 expressed in Mouse liver, Mouse testis using CWC22 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn