

LARP7 Rabbit pAb

货号: **AYP19478**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	24kDa/66kDa/67kDa
实测分子量	71kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Jurkat,Mouse testis
细胞定位	Nucleus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 160-380 of human LARP 7 (NP_057732.2).
序列	KSTGDPKGFVFEFETKEQAAKAIEFLNNPPEEAPRKPGIFPKTVKNKPIPALRVVEEKKKKKKKKGRMKKEDNIQAKEENM DTSNTSISKMKRSRPTSEGSDIESTEPQKQCSKKKKRDRVEASSLPEVRTGKRKRSSSEDAESLAPRSKVKKIIQKDIIKEA SEASKENRDIEISTEEKDTGDLKDSLLKTKRKHKKKHKERHKMGEEVIPLRVLS

靶点信息

研究背景	This gene encodes a protein which is found in the 7SK snRNP (small nuclear ribonucleoprotein). This snRNP complex inhibits a cyclin-dependent kinase, positive transcription elongation factor b, which is required for paused RNA polymerase II at a promoter to begin transcription elongation. A pseudogene of this gene is located on chromosome 3. Alternative splicing results in multiple transcript variants.
基因ID	51574

基因名	LARP7
Swiss	Q4G0J3
别名	LARP7;ALAZS;HDCMA18P;PIP7S

产品验证



Western blot analysis of LARP7 expressed in U-87MG, Jurkat, Mouse testis using LARP7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn