

— ABLYBIO, Help Your Research



# IFT88 Rabbit pAb

货号: **AYP19494**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	92kDa/93kDa/94kDa
实测分子量	108kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,BxPC-3,Mouse brain,Mouse eye
细胞定位	Cytoplasm,centriole,centrosome,cilium basal body,cytoskeleton,microtubule organizing center
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 534-833 of human IFT88 (NP_783195.2).
------	---

## 靶点信息

研究背景	This gene encodes a member of the tetratrico peptide repeat (TPR) family. Mutations of a similar gene in mouse can cause polycystic kidney disease. Two transcript variants encoding distinct isoforms have been identified for this gene.
基因ID	8100
基因名	IFT88
Swiss	Q13099 ( <a href="https://www.uniprot.org/uniprotkb/Q13099/entry">https://www.uniprot.org/uniprotkb/Q13099/entry</a> )
别名	IFT88,D13S1056E,DAF19,TG737,TTC10,hTg737,IFT88 Rabbit pAb,Recessive polycystic kidney disease protein Tg737 homolog,Tetratricopeptide repeat protein 10

## 产品验证



Western blot analysis of IFT88 expressed in Mouse brain, HepG2, BxPC-3 using IFT88 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))