

SAG Rabbit pAb

货号: **AYP19499**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	45kDa
实测分子量	52KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse eye,Rat eye
细胞定位	cytosol,photoreceptor inner segment,photoreceptor outer segment
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-405 of human SAG (NP_000532.2).
序列	MAASGKTSKSEPNHVIFKKISRDKSVTIYLGNRDYIDHVSQVQPVDPGVVLDVDPDLVKGKKVYVTLTCAFRYQGEDIDVIGLTFRRDLYFSRVQVYPPVGAASTPTKLQESLLKKGNSNTYPFLLTFPDYLPQSVMLQPAPQDSGKSCGVDFEVKAFATDSTD AEEDKIPKSSVRLLRKQVHAPLEMGPQPRAEAAWQFFMSDKPLHLAVSLNKEIYFHGEPVPTVTVTNTEKTVKKIKAFV EQVANVLYSSDYVYKPVAMEEAQEKVPPNSTLTKTLTLLPLLANNRERRGIALDGKIKHEDTNLASSTIIKEGIDRTVLGILV SYQIKVKLTVSGFLGELTSSSEVATEVPFRMLMHPQPEDPAKESYQDANLVFEEFARHNLKDAGEAEEGKRDKNDVDE

靶点信息

研究背景	Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. S-arrestin, also known as S-antigen, is a major soluble photoreceptor protein that is involved in desensitization of the photoactivated transduction cascade. It is expressed in the retina and the pineal gland and inhibits coupling of rhodopsin to transducin in vitro. Additionally, S-arrestin is highly antigenic, and is capable of inducing experimental autoimmune uveoretinitis. Mutations in this gene have been associated with Oguchi disease, a rare autosomal recessive form of night blindness.
基因ID	6295
基因名	SAG
Swiss	P10523
别名	SAG;RP47;S-AG

产品验证



Western blot analysis of SAG expressed in Mouse eye,Rat eye using SAG Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn