

— ABLYBIO, Help Your Research



MAPK12 Rabbit pAb

货号: **AYP19500**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:2000 - 1:5000 IHC: 1:50 - 1:200
理论分子量	40kDa/41kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,K-562,PC-12,22Rv1,Mouse lung,Mouse testis
细胞定位	Cytoplasm,Mitochondrion,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 248-367 of human MAP K12 (NP_002960.2).
------	---

靶点信息

研究背景	Activation of members of the mitogen-activated protein kinase family is a major mechanism for transduction of extracellular signals. Stress-activated protein kinases are one subclass of MAP kinases. The protein encoded by this gene functions as a signal transducer during differentiation of myoblasts to myotubes.
基因ID	6300
基因名	MAPK12
Swiss	P53778 (https://www.uniprot.org/uniprotkb/P53778/entry)
别名	MAPK12,ERK-6,ERK3,ERK6,MAPK 12,P38GAMMA,PRKM12,SAPK-3,SAPK3,MAPK12 Rabbit pAb,Extracellular signal-regulated kinase 6,Mitogen-activated protein kinase p38 gamma,Stress-activated protein kinase 3

产品验证



Western blot analysis of MAPK12 expressed in Mouse lung, HeLa, K-562 using MAPK12 Rabbit pAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)