

FMNL1 Rabbit pAb

货号: **AYP19507**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	121kDa/122kDa/128kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,A-549,Mouse brain
细胞定位	Cell membrane,Cell projection,Cytoplasm,Cytoplasmic vesicle,Lipid-anchor,bleb,cell cortex,phagosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 900-1070 of human FMNL1 (NP_005883.2).
序列	DLHFLDKAGSVSLDSVLADVRSLSRGLRLELTQREFVVRQDDCMVLKEFLRANSPTMDKLLADSKTAQEAFESVVEYFGENPKTTSPGLFFSLFSRFIKAYKKAQEVEQWKKEAAAQEAGADTPGKGEPPAPKSPPKARRPQMDLISELKRRQKKEPLIYESDRDGAIEDIIT

靶点信息

研究背景	This gene encodes a formin-related protein. Formin-related proteins have been implicated in morphogenesis, cytokinesis, and cell polarity. An alternative splice variant has been described but its full length sequence has not been determined.
基因ID	752

基因名	FMNL1
Swiss	O95466
别名	FMNL1;C17orf1;C17orf1B;FHOD4;FMNL;KW-13

产品验证



Western blot analysis of FMNL1 expressed in LO2,A-549,Mouse brain using FMNL1 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn