

— ABLYBIO, Help Your Research



PGK2 Rabbit pAb

货号: **AYP19522**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	44kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Jurkat,HeLa,HepG2,Mouse testis,Mouse brain,Mouse heart,Rat testis,Rat brain
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 242-360 of human PGK2 (NP_620061.2).
------	--

靶点信息

研究背景	This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of spermatogenesis.
基因ID	5232
基因名	PGK2
Swiss	P07205 (https://www.uniprot.org/uniprotkb/P07205/entry)
别名	PGK2,HEL-S-272,PGKB,PGKPS,dj417L20.2,PGK2 Rabbit pAb,Phosphoglycerate kinase,testis specific

产品验证



Western blot analysis of PGK2 expressed in Mouse testis,Rat testis,U-87MG using PGK2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)