

# PARP8 Rabbit pAb

货号: AYP19550

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	91kDa/95kDa
实测分子量	96kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse skeletal muscle,Rat skeletal muscle
细胞定位	
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human PARP8 (NP_001171527.1).
序列	MGMCSRQERIQKIDIVVIQKSRAEKDCLFADFRYSDSTFTFTYVGGPRSVSYSVHVSSEDYPDNTYVSSSENDEDVLTTE PIPVIFHRIATELRKTNDINCCLSIKSKLQKENGEEESRQNSTVEEDSEGDNDSEEFYGGQVNYDGELHKHPQLEADLSAVR EIYGPHAVSLREYGAIDDVDIDLHIDVSLDDEEIAVAWEVIRTEPIIIVRLHCSLTQYLNGPVPTVDVDFQISTKERFGLGHQLKKI MQTFVTQQWQKSKEKSNCLHNKKLSEKKVKSPLHLFSTLRRSPSYPPPGCGKS

## 靶点信息

研究背景	Mono-ADP-ribosyltransferase that mediates mono-ADP-ribosylation of target proteins.
基因ID	79668

基因名	PARP8
Swiss	Q8N3A8
别名	PARP8;ARTD16;pART16

## 产品验证



Western blot analysis of PARP8 expressed in Mouse skeletal muscle, Rat skeletal muscle using PARP8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)