

# PTGDR Rabbit pAb

货号: **AYP19558**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	32kDa/40kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 220-359 of human PTGDR (NP_000944.1).
序列	GAMRNLYAMHRRLLQRHPRSTRDCAEPRADGREASQPQLELDHLLLLALMTVLFTMCSLPVIYRAYYGAFKDVKEKNRTS EEAEDLRALRFLSVISIVDPWIFIIFRSPVFRIFHFKIFIRPLRYRSRCSNSTNMESSL

## 靶点信息

研究背景	This gene encodes a member of the guanine nucleotide-binding protein (G protein)-coupled receptor (GPCR) superfamily. The receptors are seven-pass transmembrane proteins that respond to extracellular cues and activate intracellular signal transduction pathways. This protein is reported to be a receptor for prostaglandin D2, which is a mediator of allergic inflammation and allergic airway inflammation in asthma. Alternative splicing results in multiple transcript variants.
基因ID	5729

基因名	PTGDR
Swiss	Q13258
别名	PTGDR;AS1;ASRT1;DP;DP1;PTGDR1

## 产品验证



Western blot analysis of PTGDR expressed in NIH/3T3 using PTGDR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)