

GBA2 Rabbit pAb

货号: **AYP19580**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	72kDa/98kDa/104kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse kidney,Mouse heart
细胞定位	Cytoplasmic side,Endoplasmic reticulum membrane,Golgi apparatus membrane,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 708-927 of human GBA 2 (NP_065995.1).
序列	AQDIQDKFSSILSRGQEAYERLLWNGRYNYDSSSRPQRSVMSDQCAGQWFLKACGLGEGDTEVFPTQHVVRLQTIF ELNVQAFAGGAMGAVNGMQPHGVDPKSSVQSDEVVVGVVYGLAATMIQEGLTWEGFQTAEGCYRTVWERLGLAFQT PEAYCQQRVFRSLAYMRPLSIWAMQLALQQQHKKASWPKVKQGTGLRTGPMFGPKEAMANLSPE

靶点信息

研究背景	This gene encodes a microsomal beta-glucosidase that catalyzes the hydrolysis of bile acid 3-O-glucosides as endogenous compounds. Studies to determine subcellular localization of this protein in the liver indicated that the enzyme was mainly enriched in the microsomal fraction where it appeared to be confined to the endoplasmic reticulum. This putative transmembrane protein is thought to play a role in carbohydrate transport and metabolism.
------	---

基因ID	57704
基因名	GBA2
Swiss	Q9HCG7
别名	GBA2;AD035;NLGase;SPG46

产品验证



Western blot analysis of GBA2 expressed in U-87MG, Mouse kidney, Mouse heart using GBA2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn