

— ABLYBIO, Help Your Research



VEGFR3/FLT4 Rabbit pAb

货号: **AYP19753**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	93kDa/145kDa/152kDa
实测分子量	180kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	LO2, A-549
细胞定位	Cell membrane, Cytoplasm, Nucleus, Secreted, Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1200-1300 of human VEGFR3/FLT4 (NP_891555.2).
------	--

靶点信息

研究背景	This gene encodes a tyrosine kinase receptor for vascular endothelial growth factors C and D. The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA.
基因ID	2324
基因名	FLT4
Swiss	P35916 (https://www.uniprot.org/uniprotkb/P35916/entry)
别名	FLT4,FLT-4,FLT41,LMPH1A,PCL,VEGFR-3,VEGFR3,VEGFR3/FLT4 Rabbit pAb,Fms-like tyrosine kinase 4,Tyrosine-protein kinase receptor FLT4

产品验证



Western blot analysis of VEGFR3/FLT4 expressed in LO2, A-549 using VEGFR3/FLT4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)