

# PADI2 Rabbit pAb

货号: **AYP19779**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	49kDa/75kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG, HT-29, A-549
细胞定位	Cytoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human PADI2 (NP_031391.2).
序列	MLRERTVRLQYGSRVEAVYVLGTYLWTDVYSAAPAGAQTFLKHSEHVWVEVVRDGEAEVATNGKQRWLLSPSTTLRV TMSQASTEASSDKVTNYYDEEGSIPIDQAGLFLTAIEISLDVDADRDGVVEKNNPKASWTWGPEGQGAILLVNCDRET PWLPKEDCRDEKVYSKEDLKDMSQMILRTKGPDRLPAGYEI

## 靶点信息

研究背景	This gene encodes a member of the peptidyl arginine deiminase family of enzymes, which catalyze the post-translational deimination of proteins by converting arginine residues into citrullines in the presence of calcium ions. The family members have distinct substrate specificities and tissue-specific expression patterns. The type II enzyme is the most widely expressed family member. Known substrates for this enzyme include myelin basic protein in the central nervous system and vimentin in skeletal muscle and macrophages. This enzyme is thought to play a role in the onset and progression of neurodegenerative human disorders, including Alzheimer disease and multiple sclerosis, and it has also been implicated in glaucoma pathogenesis. This gene exists in a cluster with four other paralogous genes.
基因ID	11240
基因名	PADI2
Swiss	Q9Y2J8
别名	PADI2;PAD-H19;PAD2;PDI2

## 产品验证



Western blot analysis of PADI2 expressed in U-87MG, HT-29, A-549 using PADI2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)