

— ABLYBIO, Help Your Research



CoA synthase (COASY) Rabbit pAb

货号: **AYP19802**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	62kDa/65kDa
实测分子量	62kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,293T,A-549,SKOV3,Mouse heart,Mouse liver,Rat kidney
细胞定位	Cytoplasm,Mitochondrion matrix
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 345-564 of human CoA synthase (COASY) (NP_079509.5).
------	--

靶点信息

研究背景	Coenzyme A (CoA) functions as a carrier of acetyl and acyl groups in cells and thus plays an important role in numerous synthetic and degradative metabolic pathways in all organisms. In eukaryotes, CoA and its derivatives are also involved in membrane trafficking and signal transduction. This gene encodes the bifunctional protein coenzyme A synthase (CoAsy) which carries out the last two steps in the biosynthesis of CoA from pantothenic acid (vitamin B5). The phosphopantetheine adenyltransferase domain of this bifunctional protein catalyzes the conversion of 4'-phosphopantetheine into dephospho-coenzyme A (dpCoA) while its dephospho-CoA kinase domain completes the final step by phosphorylating dpCoA to form CoA. Mutations in this gene are associated with neurodegeneration with brain iron accumulation (NBIA). Alternative splicing results in multiple isoforms.
基因ID	80347
基因名	COASY
Swiss	Q13057 (https://www.uniprot.org/uniprotkb/Q13057/entry)
别名	COASY,DPCK,NBIA6,NBP,PPAT,UKR1,CoA synthase (COASY) Rabbit pAb,POV-2,Dephospho-CoA pyrophosphorylase,Pantetheine-phosphate adenyltransferase,Dephosphocoenzyme A kinase

产品验证



Western blot analysis of CoA synthase (COASY) expressed in Mouse heart, Rat kidney, LO2 using CoA synthase (COASY) Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)