

PPIH Rabbit pAb

货号: **AYP19804**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	14kDa/19kDa
实测分子量	19kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,A-549,HeLa,Mouse testis,Rat testis
细胞定位	Cytoplasm,Nucleus speckle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-177 of human PPIH (NP_006338.1).
序列	MAVANSSPVNPVVFDDVSIQGGQEVGRMKIELFADVVPKTAENFRQFCTGFRKDGVPVIGYKGFSTFHRVIKDFMIQGGDFV NGDGTGVASIYRGPFADEFKLRHSAPGLLSMANSRGPSTNGCQFFITCSKCDWLDGKHVVFGKIIDGLVMRKIENVPTG PNNKPKLPVVISQCGEM

靶点信息

研究背景	The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is a specific component of the complex that includes pre-mRNA processing factors PRPF3, PRPF4, and PRPF18, as well as U4/U5/U6 tri-snRNP. This protein has been shown to possess PPIase activity and may act as a protein chaperone that mediates the interactions between different proteins inside the spliceosome.
------	---

基因ID	10465
基因名	PPIH
Swiss	O43447
别名	PPIH;CYP-20;CYPH;SnuCyp-20;USA-CYP

产品验证



Western blot analysis of PPIH expressed in Mouse testis,Rat testis,HepG2 using PPIH Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn