

BHMT Rabbit pAb

货号: **AYP19809**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	44kDa
实测分子量	45KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Rat liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 355-406 of human BHM T (NP_001704.2).
序列	LRIASGRPYNPSMSKPDGWSVTKGTAELMQQKEATTEQQLKELFEKQKFKSQ

靶点信息

研究背景	This gene encodes a cytosolic enzyme that catalyzes the conversion of betaine and homocysteine to dim ethylglycine and methionine, respectively. Defects in this gene could lead to hyperhomocyst(e)inemia, but such a defect has not yet been observed.
基因ID	635

基因名	BHMT
Swiss	Q93088
别名	BHMT;BHMT1;HEL-S-61p

产品验证



Western blot analysis of BHMT expressed in HepG2, Rat liver using BHMT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn