

— ABLYBIO, Help Your Research



CHD9 Rabbit pAb

货号: **AYP19815**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	324kDa/326kDa
实测分子量	300kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HeLa,Jurkat,A-549,293T,K-562,Mouse kidney
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 85-190 of human CHD9 (NP_079410.4).
------	---

靶点信息

研究背景	Acts as a transcriptional coactivator for PPARA and possibly other nuclear receptors. Proposed to be a ATP-dependent chromatin remodeling protein. Has DNA-dependent ATPase activity and binds to A/T-rich DNA. Associates with A/T-rich regulatory regions in promoters of genes that participate in the differentiation of progenitors during osteogenesis (By similarity).
基因ID	80205
基因名	CHD9
Swiss	Q3L8U1 (https://www.uniprot.org/uniprotkb/Q3L8U1/entry)
别名	CHD9,AD013,CHD-9,CReMM,KISH2,PRIC320,CHD9 Rabbit pAb,Chromatin-related mesenchymal modulator,Chromatin-remodeling factor CHROM1,Chromo domain-containing protein 9,Kismet homolog 2,PPAR-alpha-interacting complex protein 320 kDa

产品验证



Western blot analysis of CHD9 expressed in Mouse kidney,U-87MG,HeLa using CHD9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)