

DNAJC10 Rabbit pAb

货号: **AYP19817**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 37kDa/86kDa/91kDa |
| 实测分子量 | 91kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-87MG,HT-29,SKOV3,Raji,SGC-7901 |
| 细胞定位 | Endoplasmic reticulum lumen |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 614-793 of human DNAJC10 (NP_061854.1). |
| 序列 | IDCQQYHSFCAQENVQRYPEIRFFPPKSNKAYHYHSYNGWNRDAYSLRIWGLGFLPQVSTDLTPTQTFSEKVLQGNHWVI DFYAPWCGPCQNFAPEFELLARMIKGVKAGKVDCQAYAQCQKAGIRAYPTVKFYFYERAKRNFQEEQINTRDAKAI AALI SEKLETLRNQGKRKDEL |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes an endoplasmic reticulum co-chaperone which is part of the endoplasmic reticulum-associated degradation complex involved in recognizing and degrading misfolded proteins. The encoded protein reduces incorrect disulfide bonds in misfolded glycoproteins. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. |
|------|--|

| | |
|-------|--------------------------------|
| 基因ID | 54431 |
| 基因名 | DNAJC10 |
| Swiss | Q8IXB1 |
| 别名 | DNAJC10;ERdj5;JPDI;MTHr;PDIA19 |

产品验证



Western blot analysis of DNAJC10 expressed in U-87MG, HT-29, SKOV3 using DNAJC10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn