

— ABLYBIO, Help Your Research



DNAJC10 Rabbit pAb

货号: **AYP19817**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	37kDa/86kDa/91kDa
实测分子量	91kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HT-29,SKOV3,Raji,SGC-7901
细胞定位	Endoplasmic reticulum lumen
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 614-793 of human DNAJC10 (NP_061854.1).
------	---

靶点信息

研究背景	This gene encodes an endoplasmic reticulum co-chaperone which is part of the endoplasmic reticulum-associated degradation complex involved in recognizing and degrading misfolded proteins. The encoded protein reduces incorrect disulfide bonds in misfolded glycoproteins. Alternatively spliced transcript variant s encoding multiple isoforms have been observed for this gene.
基因ID	54431
基因名	DNAJC10
Swiss	Q8IXB1 (https://www.uniprot.org/uniprotkb/Q8IXB1/entry)
别名	DNAJC10,ERdj5,JPDI,MTHr,PDIA19,DNAJC10 Rabbit pAb,Dnaj homolog subfamily C member 10,Endoplasmic reticulum DNA J domain-containing protein 5,Macrothioredoxin

产品验证



Western blot analysis of DNAJC10 expressed in U-87MG, HT-29, SKOV3 using DNAJC10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)