

— ABLYBIO, Help Your Research



EIF4H Rabbit pAb

货号: **AYP19890**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	25kDa/27kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Jurkat,HeLa,SW480,Mouse pancreas
细胞定位	Cytoplasm,perinuclear region
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-228 of human EIF4H (NP_114381.1).
------	---

靶点信息

研究背景	This gene encodes one of the translation initiation factors, which functions to stimulate the initiation of protein synthesis at the level of mRNA utilization. This gene is deleted in Williams syndrome, a multisystem developmental disorder caused by the deletion of contiguous genes at 7q11.23. Alternative splicing of this gene generates 2 transcript variants.
基因ID	7458
基因名	EIF4H
Swiss	Q15056 (https://www.uniprot.org/uniprotkb/Q15056/entry)
别名	EIF4H,WBSCR1,WSCR1,eIF-4H,EIF4H Rabbit pAb,Williams-Beuren syndrome chromosomal region 1 protein,KIAA0038

产品验证



Western blot analysis of EIF4H expressed in Mouse pancreas,SH-SY5Y,Jurkat using EIF4H Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)