

NACC1 Rabbit pAb

货号: **AYP19912**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Hela,Jurkat,Mouse brain,Mouse kidney
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 170-340 of human NAC C1 (NP_443108.1).
序列	QQESDSVQCMPVAKRLWDSGQKEAGGGGNGSRKMAKFSTPDLAANRPHQPPPPQAPVVAAAQPAVAAGAGQPAGG VAAAGGVVSGPSTERTSPGTSSAYTSDSPGSYHNEEDEEEDGGEEGMDEQYRQICNMYTMYSMNVGQTAEKVEALP EQVAPESRNRIRVRQDL

靶点信息

研究背景	This gene encodes a member of the BTB/POZ protein family. BTB/POZ proteins are involved in several cellular processes including proliferation, apoptosis and transcription regulation. The encoded protein is a transcriptional repressor that plays a role in stem cell self-renewal and pluripotency maintenance. The encoded protein also suppresses transcription of the candidate tumor suppressor Gadd45GIP1, and expression of this gene may play a role in the progression of multiple types of cancer. A pseudogene of this gene is located on the short arm of chromosome 9.
------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	112939
基因名	NACC1
Swiss	Q96RE7
别名	NACC1;BEND8;BTBD14B;BTBD30;NAC-1;NAC1;NECFM

产品验证



Western blot analysis of NACC1 expressed in Mouse brain,SKOV3,Hela using NACC1 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn