

— ABLYBIO, Help Your Research



SH2D1A Rabbit pAb

货号: **AYP19970**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	6kDa/8kDa/13kDa/14kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,PC-3,Mouse spleen
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-128 of human SH2D1 A (NP_002342.1).
------	---

靶点信息

研究背景	This gene encodes a protein that plays a major role in the bidirectional stimulation of T and B cells. This protein contains an SH2 domain and a short tail. It associates with the signaling lymphocyte-activation molecule, thereby acting as an inhibitor of this transmembrane protein by blocking the recruitment of the SH2-domain-containing signal-transduction molecule SHP-2 to its docking site. This protein can also bind to other related surface molecules that are expressed on activated T, B and NK cells, thereby modifying signal transduction pathways in these cells. Mutations in this gene cause lymphoproliferative syndrome X-linked type 1 or Duncan disease, a rare immunodeficiency characterized by extreme susceptibility to infection with Epstein-Barr virus, with symptoms including severe mononucleosis and malignant lymphoma. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	4068
基因名	SH2D1A
Swiss	O60880 (https://www.uniprot.org/uniprotkb/O60880/entry)
别名	SH2D1A,DSHP,EBVS,IMD5,LYP,MTCP1,SAP,SAP/SH2D1A,XLP,XLPD,XLPD1,SH2D1A Rabbit pAb,Duncan disease SH2-protein,Signaling lymphocytic activation molecule-associated protein,T-cell signal transduction molecule SAP

产品验证



Western blot analysis of SH2D1A expressed in Jurkat,PC-3,Mouse spleen using SH2D1A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)