

TRRAP Rabbit pAb

货号: **AYP19980**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	
实测分子量	434kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Rat testis
细胞定位	Golgi apparatus,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 3660-3859 of TRRAP (NP_001231509.1).
序列	PNATDYWTFRKMFTIQLALIGFAEFVLHLNRLNPEMLQIAQDTGKLNVAIFYFRFDINDATGDLNANRPVFPRLTPNISEFLTTI GVSGPLTASMIAVARCFAQPNFKVDGILKTVLRDEIIAWHKKTQEDTSSPLSAAGQPENMDSQQLVSLVQKAVTAIMTRLH NLAQFEGGESKVNTLVAAANSLDNLRCRMDPAWHPWL

靶点信息

研究背景	This gene encodes a large multidomain protein of the phosphoinositide 3-kinase-related kinases (PIKK) family. The encoded protein is a common component of many histone acetyltransferase (HAT) complexes and plays a role in transcription and DNA repair by recruiting HAT complexes to chromatin. Deregulation of this gene may play a role in several types of cancer including glioblastoma multiforme. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Sep 2011]
------	--

基因ID	8295
基因名	TRRAP
Swiss	Q9Y4A5
别名	PAF350/400; PAF400; STAF40; TR-AP; Tra1

产品验证



Western blot analysis of TRRAP expressed in HeLa, Rat testis using TRRAP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn