

— ABLYBIO, Help Your Research



TRRAP Rabbit pAb

货号: **AYP19980**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	
实测分子量	434kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Rat testis
细胞定位	Golgi apparatus,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 3660-3859 of TRRAP (N P_001231509.1).
------	---

靶点信息

研究背景	This gene encodes a large multidomain protein of the phosphoinositide 3-kinase-related kinases (PIKK) family. The encoded protein is a common component of many histone acetyltransferase (HAT) complexes and plays a role in transcription and DNA repair by recruiting HAT complexes to chromatin. Deregulation of this gene may play a role in several types of cancer including glioblastoma multiforme. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Sep 2011]
基因ID	8295
基因名	TRRAP
Swiss	Q9Y4A5 (https://www.uniprot.org/uniprotkb/Q9Y4A5/entry)
别名	PAF350/400,PAF400,STAF40,TR-AP,Tra1,TRRAP Rabbit pAb,TRRAP,350/400 kDa PCAF-associated factor,Tra 1 homolog

产品验证



Western blot analysis of TRRAP expressed in HeLa,Rat testis using TRRAP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)