

NCOA1 Rabbit pAb

货号: **AYP19981**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	152kDa/156kDa
实测分子量	157kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,K-562
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 720-980 of human NCOA1 (NP_003734.3).
序列	SVTGQVQGNSSIKLELDASKKKESKDHQLLRYLLDKDEKDLRSTPNLSLDDVKVKVEKKEQMDPCNTNPTPMTKPTPEEKLEAQSQFTADLDQFDQLLPTLEKAAQLPGLCETDRMDGAVTSVTIKSEILPASLQSATARPTSRLNRLPELELEAIDNQFGQPGTGDQIPWTNNTVTAINQSKSEDQCISSQLDELLCPPTTVEGRNDEKALLEQLVSFSLSGKDETELAELDRALGIDKLVQGGGLDVLSERFPPQATP

靶点信息

研究背景	The protein encoded by this gene acts as a transcriptional coactivator for steroid and nuclear hormone receptors. It is a member of the p160/steroid receptor coactivator (SRC) family and like other family members has histone acetyltransferase activity and contains a nuclear localization signal, as well as bHLH and PAS domains. The product of this gene binds nuclear receptors directly and stimulates the transcriptional activities in a hormone-dependent fashion. Alternatively spliced transcript variants encoding different isoforms have been identified.
------	--

基因ID	8648
基因名	NCOA1
Swiss	Q15788
别名	NCOA1;F-SRC-1;KAT13A;RIP160;SRC1;bHLHe42;bHLHe74

产品验证



Western blot analysis of NCOA1 expressed in NIH/3T3, K-562 using NCOA1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn