

C19orf10 Rabbit pAb

货号: **AYP20007**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:500
理论分子量	18kDa
实测分子量	19kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-431,Mouse liver,Rat pancreas
细胞定位	Endoplasmic reticulum-Golgi intermediate compartment,Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-173 of human C19orf10 (NP_061980.1).
序列	MAAPSGGWNGVGASLWAALLLGAVALRPAEAVSEPTTVAFDVVRPGGVVHSHNVGPGDKYTCMFTYASQGGTNEQWQMSLGTSEDHQHFTCTIWRPQGKSYLYFTQKAEVRGAEIEYAMAYSKAAFERESDVPLKTEEFVTKTAVAHRPGAFKAE LSKLVIVAKASRTEL

靶点信息

研究背景	The protein encoded by this gene was previously thought to support proliferation of lymphoid cells and was considered an interleukin. However, this activity has not been reproducible and the function of this protein is currently unknown.
基因ID	56005

基因名	MYDGF
Swiss	Q969H8
别名	MYDGF;C19orf10;EUROIMAGE1875335;IL25;IL27;IL27w;R33729_1;SF20

产品验证



Western blot analysis of C19orf10 expressed in Mouse liver,Rat pancreas,293T using C19orf10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn