

UCK2 Rabbit pAb

货号: **AYP20018**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	12kDa/29kDa
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HeLa,293T,B-cell,Mouse thymus
细胞定位	cytoplasm,cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-261 of human UCK2 (NP_036606.2).
序列	MAGDSEQTLQNHQQPNNGGEPFLIGVSGGTASGKSSVCAKIVQLLQNEVDYRQKQVWILSQDSFYRVLTSQKAKALKG QFNFDHPDAFDNELILKTLKEITEGKTVQIPVYDFVSHSRKEETVTVYPADVVLFEGLAFYSQEVRLDFQMCLFVDTDADT RLSRRVLRDISERGRDLEQILSQYITFVKPAFEFCLPTKKYADVIIIPRGADNLVAINLIVQHIQDILNGGPSKRQTNGCLNGYT PSRKRQASESSSRPH

靶点信息

研究背景	This gene encodes a pyrimidine ribonucleoside kinase. The encoded protein (EC 2.7.1.48) catalyzes phosphorylation of uridine and cytidine to uridine monophosphate (UMP) and cytidine monophosphate (CMP), respectively.
基因ID	7371

基因名	UCK2
Swiss	Q9BZX2
别名	UCK2;TSA903;UK;UMPK

产品验证



Western blot analysis of UCK2 expressed in Mouse thymus,A-549,HeLa using UCK2 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn