

WHRN Rabbit pAb

货号: **AYP20026**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	37kDa/55kDa/59kDa/96kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HeLa,K-562,Mouse liver
细胞定位	Cell projection,Cytoplasm,growth cone,sterocilium
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 708-907 of human WHR N (NP_056219.3).
序列	PSGHPDQTGTNQHFVMVEVHRPDPSEPDVNEVRALPQTRTASTLSQLSDSGQTLSEDSGVDAGEAEASAPGRGRQSVST KSRSSKELPRNERPTDGANKPPGLLEPTSTLVRVKKSAATLGIAIEGGANTRQPLPRIVTIQRGGSAHNCGQLKVGHVILEV NGLTLRGKEHREAARIIEAFKTKDRDYIDFLVTEFNVML

靶点信息

研究背景	This gene is thought to function in the organization and stabilization of sterocilia elongation and actin cytoskeletal assembly, based on studies of the related mouse gene. Mutations in this gene have been associated with autosomal recessive non-syndromic deafness and Usher Syndrome. Alternative splicing results in multiple transcript variants.
基因ID	25861

基因名	WHRN
Swiss	Q9P202
别名	WHRN;CIP98;DFNB31;PDZD7B;USH2D;Wl;whirlin

产品验证



Western blot analysis of WHRN expressed in Mouse liver,U-87MG,HeLa using WHRN Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn