

— ABLYBIO, Help Your Research



# TNFRSF17 Rabbit pAb

货号: **AYP20034**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	14kDa/20kDa
实测分子量	15-35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SP2/0,Raji,Rat spleen
细胞定位	Cell membrane,Endomembrane system,Single-pass type III membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human TNFRSF17 (NP_001183.2).
------	---

## 靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation.
基因ID	608
基因名	TNFRSF17
Swiss	Q02223 ( <a href="https://www.uniprot.org/uniprotkb/Q02223/entry">https://www.uniprot.org/uniprotkb/Q02223/entry</a> )
别名	TNFRSF17,BCM,BCMA,CD269,TNFRSF13A,TNFRSF17 Rabbit pAb,B-cell maturation protein

## 产品验证



Western blot analysis of TNFRSF17 expressed in SP2/0,Raji,Rat spleen using TNFRSF17 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)