

— ABLYBIO, Help Your Research



TDGF1 Rabbit pAb

货号: **AYP20043**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	21kDa
实测分子量	18KDa/20KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NCCIT
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 15-134 of human TDGF 1 (NP_001167607.1).
------	--

靶点信息

研究背景	This gene encodes an epidermal growth factor-related protein that contains a cripto, FRL-1, and cryptic domain. The encoded protein is an extracellular, membrane-bound signaling protein that plays an essential role in embryonic development and tumor growth. Mutations in this gene are associated with forebrain defects. Pseudogenes of this gene are found on chromosomes 2, 3, 6, 8, 19 and X. Alternate splicing results in multiple transcript variants.
基因ID	6997
基因名	TDGF1
Swiss	P13385 (https://www.uniprot.org/uniprotkb/P13385/entry)
别名	TDGF1,CR,CRGF,CRIPTO,TDGF1 Rabbit pAb,EGF-CFC family member,Cripto-1 growth factor,Epidermal growth factor-like cripto protein CR1,Teratocarcinoma-derived growth factor 1,CRIPTO-1

产品验证



Western blot analysis of TDGF1 expressed in NCCIT using TDGF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)