

JMJD8 Rabbit pAb

货号: **AYP20086**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	33kDa/36kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,HL-60,Jurkat,Mouse kidney,Mouse liver,Rat testis,Rat liver
细胞定位	cytoplasm,endoplasmic reticulum,endoplasmic reticulum lumen,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-285 of human JMJD8 (NP_001005920.2).
序列	MAGGWAQGSRRARLPEGGRLMAPASRLALWALAVALPGSGAEGDGGWRPGGPGAVAEERCTVERRADLTAEF VQYAFVRPVILQGLTDNSRFRALCSRDRLLASFGDRVRLSTANTYSYHKVDLPFQEYVEQLLHPQDPTSLGNDTLYFFG DNNFTEWASLFRHYSPPFGLGTAPAYSFGIAGAGSGVPFHWHGPGYSEVIYGRKRWFYPPPEKTPEFHPNKTTLAWLR DTYPALPPSARPLECTIRAGEVLYFPDRWWHATLNLDTSVFISTFLG

靶点信息

研究背景	Functions as a positive regulator of TNF-induced NF-kappa-B signaling. Regulates angiogenesis and cellular metabolism through interaction with PKM.
基因ID	339123

基因名	JMJD8
Swiss	Q96S16
别名	JMJD8;C16orf20;PP14397

产品验证



Western blot analysis of JMJD8 expressed in Mouse kidney,Rat testis,293T using JMJD8 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn