

ATF7IP Rabbit pAb

货号: **AYP20092**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	118kDa/136kDa/137kDa
实测分子量	250KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,A-549,Mouse kidney
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 525-725 of human ATF7 IP (NP_060649.3).
序列	NEKDNKPEEEEEQVIHEDDERPSEKNEFSRRKRKSKSEMDNVQSKRRRYMEEYEAEFQVKITAKGDINQKLQKVIQWLLLEKLCALQCAVFDKTLAELKTRVEKIECNKRHKTVLTELQAKIARLTKRFEAAKEDLKKRHEHPPNPPVSPGKTVNDVNSNNNMSYRNAGTVRQMLESKRNVSESAPPSFQTPVNTVSSTNL

靶点信息

研究背景	ATF7IP is a multifunctional nuclear protein that associates with heterochromatin. It can act as a transcriptional coactivator or corepressor depending upon its binding partners (summary by Liu et al., 2009 [PubMed 19106100]).[supplied by OMIM, Nov 2010]
基因ID	55729

基因名	ATF7IP
Swiss	Q6VMQ6
别名	ATF7IP;AM;ATF-IP;MCAF;MCAF1;p621

产品验证



Western blot analysis of ATF7IP expressed in MCF7,A-549,Mouse kidney using ATF7IP Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn