

# XPO6 Rabbit pAb

货号: **AYP20096**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	127kDa/128kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse testis,Mouse brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 290-530 of human XPO6 (NP_055986.1).
序列	ASVNGSSQNCVSGQERGLGVLAMSCINELMSKNCVPMEFEEYLLRMFQQTFYLLQKITKDNNHAHTVKSRL EELDES YIEK FTDFLRLFVSVHLRRIESYSQFPVVEFLTL LFKYTFHQPTHEGYFSCLDIWTLFLDYLT SKIKSRLGDKEAVLNRYEDALVLLL TEVLNRIQFRYNQAQLEELDDETLDDDDQQTEWQRYLRQSLEVVAKVMELLPTHAFSTLFPVLQDNLEVYLG LQQFI

## 靶点信息

研究背景	The protein encoded by this gene is a member of the importin-beta family. Members of this family are regulated by the GTPase Ran to mediate transport of cargo across the nuclear envelope. This protein has been shown to mediate nuclear export of profilin-actin complexes. A pseudogene of this gene is located on the long arm of chromosome 14. Alternative splicing results in multiple transcript variants that encode different protein isoforms.
------	--

基因ID	23214
基因名	XPO6
Swiss	Q96QU8
别名	XPO6;EXP6;RANBP20

## 产品验证



Western blot analysis of XPO6 expressed in Jurkat, Mouse testis, Mouse brain using XPO6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)