

DIAPH3 Rabbit pAb

货号: **AYP20114**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	79kDa/98kDa/127-136kDa
实测分子量	137kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	MCF7, HeLa, HT-1080, NCI-H460
细胞定位	Cytoplasm, cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 630-849 of human DIAPH3 (NP_112194.2).
序列	KYPDILNFVDDLEPLDKASKVSVETLEKNLRQMGRQLQLEKELETFPDPEDLHDKFVTKMSRFVISAKEQYETLSKLHEN MEKLYQSIIGYYAIDVKKVSVEDFLTDLNNFRRTTFMQAIKENIKKREAEKEKRVRIAKELAERERLERQKKRLLLEMKTEG DETGVMNDNLLEALQSGAAFRDRRKRTPMPKDVRQSLSPMSQRPVLKVCNHNKPYL

靶点信息

研究背景	This gene encodes a member of the diaphanous subfamily of the formin family. Members of this family are involved in actin remodeling and regulate cell movement and adhesion. Mutations in this gene are associated with autosomal dominant auditory neuropathy 1. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	81624

基因名	DIAPH3
Swiss	Q9NSV4
别名	DIAPH3;AN;AUNA1;DIA2;DRF3;NSDAN;diap3;mDia2

产品验证



Western blot analysis of DIAPH3 expressed in MCF7, HeLa, HT-1080 using DIAPH3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn