

— ABLYBIO, Help Your Research



FCRL4 Rabbit pAb

货号: **AYP20120**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	56kDa/57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,B cells,293T,Mouse kidney,Mouse liver,Rat kidney,Rat liver
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 120-387 of human FCRL 4 (NP_112572.1).
------	--

靶点信息

研究背景	This gene encodes a member of the immunoglobulin receptor superfamily and is one of several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein has four extracellular C2-type immunoglobulin domains, a transmembrane domain and a cytoplasmic domain that contains three immune-receptor tyrosine-based inhibitory motifs. This protein may play a role in the function of memory B-cells in the epithelia. Aberrations in the chromosomal region encoding this gene are associated with non-Hodgkin lymphoma and multiple myeloma.
基因ID	83417
基因名	FCRL4
Swiss	Q96PJ5 (https://www.uniprot.org/uniprotkb/Q96PJ5/entry)
别名	FCRL4,CD307d,FCRH4,IGFP2,IRTA1,FCRL4 Rabbit pAb,Fc receptor homolog 4,IFGP family protein 2,Immune receptor translocation-associated protein 1,IFGP2

产品验证



Western blot analysis of FCRL4 expressed in Mouse kidney,Rat kidney,Raji using FCRL4 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)