

# FCRL4 Rabbit pAb

货号: AYP20120

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:100 - 1:200
理论分子量	56kDa/57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,B cells,293T,Mouse kidney,Mouse liver,Rat kidney,Rat liver
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 120-387 of human FCRL4 (NP_112572.1).
序列	VLRCHRRRKEKLTAVKYTWNGNILSISNKSVDLLIPQASSNNGNYRCIGYGDENDVFRSNFKIIKIQLFPHPELKATDSQ PTEGNSVNLSCETQLPPERSDTPLHFNFRRDGEVILSDWSTYPELQLPTVWRENSGSYWCGAETVRGNIHKHSPSLQIHV QRIPVSGVLLLETQPSGGQAVEGEMLVLVCSVAEGTGDITFSWHREDMQESLGRKTQRSLRAELELPAIRQSHAGGYCT ADNSYGPVQSMVLNVTVRETPGNRDGL

## 靶点信息

研究背景	This gene encodes a member of the immunoglobulin receptor superfamily and is one of several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein has four extracellular C2-type immunoglobulin domains, a transmembrane domain and a cytoplasmic domain that contains three immune-receptor tyrosine-based inhibitory motifs. This protein may play a role in the function of memory B-cells in the epithelia. Aberrations in the chromosomal region encoding this gene are associated with non-Hodgkin lymphoma and multiple myeloma.
基因ID	83417
基因名	FCRL4
Swiss	Q96PJ5
别名	FCRL4;CD307d;FCRH4;IGFP2;IRTA1

## 产品验证



Western blot analysis of FCRL4 expressed in Mouse kidney, Rat kidney, Raji using FCRL4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)