

— ABLYBIO, Help Your Research



# SLC28A3 Rabbit pAb

货号: **AYP20121**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	69kDa/76kDa
实测分子量	77kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,293T
细胞定位	Cell membrane,Endoplasmic reticulum membrane,Multi-pass membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 612-691 of human SLC28A3 (NP_071410.1).
------	---

## 靶点信息

研究背景	Nucleoside transporters, such as SLC28A3, regulate multiple cellular processes, including neurotransmission, vascular tone, adenosine concentration in the vicinity of cell surface receptors, and transport and metabolism of nucleoside drugs. SLC28A3 shows broad specificity for pyrimidine and purine nucleosides (Ritzel et al., 2001 [PubMed 11032837]).[supplied by OMIM, Mar 2008]
基因ID	64078
基因名	SLC28A3
Swiss	Q9HAS3 ( <a href="https://www.uniprot.org/uniprotkb/Q9HAS3/entry">https://www.uniprot.org/uniprotkb/Q9HAS3/entry</a> )
别名	SLC28A3,CNT3,SLC28A3 Rabbit pAb,Concentrative Na(+)-nucleoside cotransporter 3

## 产品验证



Western blot analysis of SLC28A3 expressed in HepG2,293T using SLC28A3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))