

# LRIG1 Rabbit pAb

货号: **AYP20132**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:1000 - 1:2000
理论分子量	116kDa/119kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse skin,Mouse brain
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 816-1093 of human LRI G1 (NP_056356.2).
序列	YQTRKKSEEYSVTNTDETVPDPVPSYLSQGTLSDRQETVVRTEGGPQANGHIESNGVCPRDASHFPEPDTHSVACRQPKLCAGSAYHKEPWKAMEKAEGTPGPHKMEHGGRVVCSDCNTEVDCYSRGQAFHPQPVS RDSAQPSAPNGPEPGGSDQEHSPPHHQCSRTAAGSCPECQGSLYPSNHDRMLTAVKKKPMASLDGKGDSSWTLARLYHPDSTELQPASSLTSGSPERAEAQYLLVSNHLPKACDASPESTPLTGQLPGKQRVPLLLAPKS

## 靶点信息

研究背景	Acts as a feedback negative regulator of signaling by receptor tyrosine kinases, through a mechanism that involves enhancement of receptor ubiquitination and accelerated intracellular degradation.
基因ID	26018

基因名	LRIG1
Swiss	Q96JA1
别名	LRIG1;LIG-1;LIG1

## 产品验证



Western blot analysis of LRIG1 expressed in Raji, Mouse skin, Mouse brain using LRIG1 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)