

# BICD2 Rabbit pAb

货号: **AYP20140**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:1000 - 1:2000
理论分子量	93kDa/96kDa
实测分子量	118kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HL-60,HeLa,Mouse brain,Rat brain
细胞定位	Cytoplasm,Golgi apparatus,cytoskeleton
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 440-620 of human BICD 2 (NP_001003800.1).
序列	HVAVAEAGELREQLKALRSTHEAREAQHAEEKGRYEAEQGALTEKVSLEKASRQDRELLARLEKELKKVSDVAGETQGSL SVAQDELVTFSEELANLYHHVCMCNNETPNRVMLDYREGQGAGRTSPGGRTSPEARARRSPILLPKGLLAPEAGRAD GGTGDSPPSPGSSLPSPSP

## 靶点信息

研究背景	This gene is one of two human homologs of Drosophila bicaudal-D and a member of the Bicoid family. It has been implicated in dynein-mediated, minus end-directed motility along microtubules. It has also been reported to be a phosphorylation target of NIMA related kinase 8. Two alternative splice variants have been described.
基因ID	23299

基因名	BICD2
Swiss	Q8TD16
别名	BICD2;SMALED2;bA526D8.1

## 产品验证



Western blot analysis of BICD2 expressed in Mouse brain,Rat brain,U-87MG using BICD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)