

— ABLYBIO, Help Your Research



# Cyclin K Rabbit pAb

货号: **AYP20147**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	40kDa/41kDa/64kDa/66kDa
实测分子量	90kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse brain
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human Cyclin K (NP_001092872.1).
------	---

## 靶点信息

研究背景	The protein encoded by this gene is a member of the transcription cyclin family. These cyclins may regulate transcription through their association with and activation of cyclin-dependent kinases (CDK) that phosphorylate the C-terminal domain (CTD) of the large subunit of RNA polymerase II. This gene product may play a dual role in regulating CDK and RNA polymerase II activities.
基因ID	8812
基因名	CCNK
Swiss	O75909 ( <a href="https://www.uniprot.org/uniprotkb/O75909/entry">https://www.uniprot.org/uniprotkb/O75909/entry</a> )
别名	CCNK,CPR4,cyclin-K,Cyclin K Rabbit pAb

## 产品验证



Western blot analysis of Cyclin K expressed in HeLa, Mouse brain using Cyclin K Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)