

ZSCAN4C Rabbit pAb

货号: **AYP20168**

产品信息

| | |
|-------|--|
| 反应 | Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 57kDa |
| 实测分子量 | 57kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse kidney |
| 细胞定位 | Chromosome,Nucleus,telomere |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 400-506 of mouse ZSCAN4 (NP_001013787.1). |
| 序列 | CSRMFKHARSLSSHQRTHLNKKSELLCVTCQKMFKRVSDDRRTHEIIHMPEKPFKCSTCEKSFHKTNLKSHEMIHTGEMPYVCSLCSRRFRQSSTYHRHLRNYHRSD |

靶点信息

| | |
|------|--|
| 研究背景 | Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Acts upstream of or within negative regulation of mitotic recombination; regulation of transcription by RNA polymerase II; and telomere maintenance via telomere lengthening. Located in chromosome, telomeric region; cytoplasm; and nucleus. Is expressed in 1-cell stage embryo; 2-cell stage embryo; primary oocyte; and secondary oocyte. Orthologous to human ZSCAN4 (zinc finger and SCAN domain containing 4). |
|------|--|

| | |
|-------|-----------------------|
| 基因ID | 245109 |
| 基因名 | |
| Swiss | |
| 别名 | ZSCAN4;ZNF494;ZSCAN4C |

产品验证



Western blot analysis of ZSCAN4C expressed in Mouse kidney using ZSCAN4C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn