

PADI1 Rabbit pAb

货号: **AYP20189**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	74kDa
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse ovary,Rat uterus
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human PADI1 (NP_037490.2).
序列	MAPKRVVQLSLKMPHAVCVVGVEAHVDIHSVDPKGANSFRVSGSSGVEVFMVYNRTRVKEPIGKARWPLDTDADMVV SVGTASKELKDFKVRVSYFGEQEDQALGRSVLYLTGVDISLEVDTGRTGKVKRSQGDKKTWRWGPEGYGAILLVNCDR DNHRSAEPDLTHSWLMSLADLQDMSPMLLSCNGPDKLFDHKLVLNVPFSDSKRVRVFCARGGNSLSDYKQVLGPQCL SYEVERQPGEQEIFYVEGLTFPDAD

靶点信息

研究背景	This gene encodes a member of the peptidyl arginine deiminase family of enzymes, which catalyze the post-translational deimination of proteins by converting arginine residues into citrullines in the presence of calcium ions. The family members have distinct substrate specificities and tissue-specific expression patterns. The type I enzyme is involved in the late stages of epidermal differentiation, where it deiminates filaggrin and keratin K1, which maintains hydration of the stratum corneum, and hence the cutaneous barrier function. This enzyme may also play a role in hair follicle formation. This gene exists in a cluster with four other paralogous genes.
基因ID	29943
基因名	PADI1
Swiss	Q9ULC6
别名	PADI1;HPAD10;PAD1;PDI;PDI1

产品验证



Western blot analysis of PADI1 expressed in Mouse ovary, Rat uterus using PADI1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn