

MSMB Rabbit pAb

货号: **AYP20216**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	8kDa/12kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,293T
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-114 of human MSMB (NP_002434.1).
序列	VPGDSTRKCMDLKGKHPINSEWQTDNCETCTCYETEISCTLVSTPVGYDKDNCQRIFKKEDCKYIVVEKKDPKKTCSVSEWII

靶点信息

研究背景	The protein encoded by this gene is a member of the immunoglobulin binding factor family. It is synthesized by the epithelial cells of the prostate gland and secreted into the seminal plasma. This protein has inhibin-like activity. It may have a role as an autocrine paracrine factor in uterine, breast and other female reproductive tissues. The expression of the encoded protein is found to be decreased in prostate cancer. Two alternatively spliced transcript variants encoding different isoforms are described for this gene. The use of alternate polyadenylation sites has been found for this gene.
------	--

基因ID	4477
基因名	MSMB
Swiss	P08118
别名	MSMB;HPC13;IGBF;MSP;MSPB;PN44;PRPS;PSP;PSP-94;PSP57;PSP94

产品验证



Western blot analysis of MSMB expressed in SW480,293T using MSMB Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn