

FZD3 Rabbit pAb

货号: **AYP20231**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	76kDa
实测分子量	76KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	Apical cell membrane,Cell membrane,Cell surface,Membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 23-205 of human FZD3 (NP_059108.1).
序列	HSLFSCPEITLRMCQDLPYNTTFMPNLLNHYDQQTAAALAMEPFHPMVNLDCSRDFRPFLCALYAPICMEYGRVTLPCRRLC QRAYSECSKLMEMFGVPWPEDMECSRFPDCDEPYRRLVDLNLAGEPTEGAPVAVQRDYGFWCPRELKIDPDLGYSFLHV RDCSPPCPNMYFREELSFARYF

靶点信息

研究背景	This gene is a member of the frizzled gene family. Members of this family encode seven-transmembrane domain proteins that are receptors for the wiggless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. The function of this protein is unknown, although it may play a role in mammalian hair follicle development. Alternative splicing results in multiple transcript variants. This gene is a susceptibility locus for schizophrenia.
------	--

基因ID	7976
基因名	FZD3
Swiss	Q9NPG1
别名	FZD3;Fz-3

产品验证



Western blot analysis of FZD3 expressed in Rat brain using FZD3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn