

NOSIP Rabbit pAb

货号: **AYP20245**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	33kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,B cells,293T
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-301 of human NOSIP (NP_057037.1).
序列	MTRHGKNCTAGAVYTYHEKKKDTAASGYGTQNIIRLSRDAVKDFDCCCLSLQPCHDPVPTDGYLYEREIAILEYLHQKKEI ARQMKAYEKQGRTRREEQKELQRAASQDHVRGFLEKESAIVSRPLNPFTAKALSGTSPDDVQPGPSVGPPSKDKDKVLP SFWIPSLTPEAKATKLEKPSRTVTCMSGKPLRMSDLTPVHFTPLDSSVDRVGLITRSERYVCAVTRDLSLNATPCAVLRPS GAVVTLECVKLRKDMVDPVTGDKLTDRIIVLQRGGTGFGAGSGVKLQAEKSRPVMQA

靶点信息

研究背景	The protein encoded by this gene may modulate the activity and localization of nitric oxide synthase (endothelial and neuronal) and thus nitric oxide production. Alternative splicing results in multiple transcript variants that encode the same protein.
基因ID	51070

基因名	NOSIP
Swiss	Q9Y314
别名	NOSIP;CGI-25

产品验证



Western blot analysis of NOSIP expressed in A-549,B cells,293T using NOSIP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn