

— ABLYBIO, Help Your Research



# SNX9 Rabbit pAb

货号: **AYP20262**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 | 66kDa  |
| 实测分子量 | 90kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.   |
| 偶联物   | Unconjugated   |
| 阳性对照  | A375,HeLa,Mouse liver,Rat liver,Rat heart  |
| 细胞定位  | Cell membrane,Cell projection,Cytoplasm,Cytoplasmic side,Cytoplasmic vesicle,Cytoplasmic vesicle membrane,Golgi apparatus,Peripheral membrane protein,clathrin-coated vesicle,ruffle,trans-Golgi network |
| 纯化    | Affinity purification  |


## 抗原信息

|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-210 of human SNX9 (NP_057308.1). |
|------|--|

## 靶点信息

|       |   |
|-------|---|
| 研究背景  | This gene encodes a member of the sorting nexin family. Members of this family contain a phosphoinositide binding domain, and are involved in intracellular trafficking. The encoded protein does not contain a coiled coil region, like some family members, but does contain a SRC homology domain near its N-terminus. The encoded protein is reported to have a variety of interaction partners, including of adaptor protein 2, dynamin, tyrosine kinase non-receptor 2, Wiskott-Aldrich syndrome-like, and ARP3 actin-related protein 3. The encoded protein is implicated in several stages of intracellular trafficking, including endocytosis, macropinocytosis, and F-actin nucleation. |
| 基因ID  | 51429   |
| 基因名   | SNX9  |
| Swiss | Q9Y5X1 ( <a href="https://www.uniprot.org/uniprotkb/Q9Y5X1/entry">https://www.uniprot.org/uniprotkb/Q9Y5X1/entry</a> )  |
| 别名    | SNX9,SDP1,SH3PX1,SH3PXD3A,WISP,SNX9 Rabbit pAb,SH3 and PX domain-containing protein 1,SH3 and PX domain-containing protein 3A   |

## 产品验证



Western blot analysis of SNX9 expressed in Mouse liver,Rat liver,A375 using SNX9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)