

# HDAC4 Rabbit pAb

货号: AYP20328

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:100 <b>IF/ICC:</b> 1:100 - 1:200
理论分子量	106kDa/119kDa
实测分子量	140KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C2C12
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 451-650 of human HDA C4 (NP_006028.2).
序列	PSIHKLRQHRPLGRTQSAPLPQNAQALQHLVIQQHQFLEKHKQQFQQQLQMNKIIPKSEPARQPESHPEETEEELRE HQALLDEPYLDRLPGQKEAHAQAGVQVKQEPIDESDEEEAEPREVEPGQRQPSEQELLFRQQALLLEQQRIHQLRNYQAS MEAAGIPVSGGHRPLSRAQSSPASATFPVSVQEPPTKP

## 靶点信息

研究背景	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events . Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.
基因ID	9759
基因名	HDAC4
Swiss	P56524
别名	HDAC4;AHO3;BDMR;HA6116;HD4;HDAC-4;HDAC-A;HDACA

## 产品验证



Western blot analysis of HDAC4 expressed in C2C12 using HDAC4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)