

— ABLYBIO, Help Your Research



Phospho-PEA15-S104 Rabbit pAb

货号: **AYP20378**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	15kDa/17kDa
实测分子量	13kDa/15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,C6
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S104 of human PEA15 (NP_003759.1).
------	--

靶点信息

研究背景	This gene encodes a death effector domain-containing protein that functions as a negative regulator of a poptosis. The encoded protein is an endogenous substrate for protein kinase C. This protein is also overe xpressed in type 2 diabetes mellitus, where it may contribute to insulin resistance in glucose uptake. Alte rnative splicing results in multiple transcript variants.
基因ID	8682
基因名	PEA15
Swiss	Q15121 (https://www.uniprot.org/uniprotkb/Q15121/entry)
别名	PEA15,HMAT1,HUMMAT1H,MAT1,MAT1H,PEA-15,PED,Phospho-PEA15-S104 Rabbit pAb,15 kDa phosphopro tein enriched in astrocytes,Phosphoprotein enriched in diabetes

产品验证



Western blot analysis of Phospho-PEA15-S104 expressed in HeLa,C6 using Phospho-PEA15-S104 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per l ane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)